

Correspondence

Painted Spurfowl from the Chota Nagpur Plateau area of West Bengal

The Painted Spurfowl *Galloperdix lunulata* inhabits rocky slopes and scrub habitats of dry forests, from central to southern India, including the Aravalli Range of Rajasthan up to the Ayodhya Range of West Bengal (Ball 1874). Historically, it was recorded at Rajmahal Hill, Manbhum, Lohardaga, Sirguja, Jashpur, Udaipur, Sambalpur, northwards of Mahanadi, Raipur, Nowagarh, Karial, and along the Godavari Valley (Ball 1874, 1878) on the Chotanagpur Plateau. But there is no record of the species (Chakraborty 2011) from a large portion of this important ecogeographic region (Kumar & Rawat 2008). We report the presence of this species in the Chotanagpur region on the basis of several sight records supported by photographs taken on three occasions [163].



163. Male Painted Spurfowl.

Ayan Mondal and Anirban Patra first saw the bird in the Matha forest, Purulia District on 16 April 2013. AP and Anirban Patra photographed a bird on 15 November 2015 in the Jhilimili forest (22.84°N, 86.71°E). Subsequently, SS and Kirty Kumar photographed a cock on 05 February 2019 on Ajodhya Hill (23.20°N, 86.07°E). DS and SM saw a pair on 28 April 2019 on the outskirts of Bishnupur town (23.03°E, 87.29°N). This was typical plateau habitat (Fig. 1), with its characteristic red soil and mounds of granite rock, with scarce vegetation like date palm, sal, palas, eucalyptus, etc. We did not see a waterbody.

According to Grimm et al. (2011), although this bird is found on the Chotanagpur Plateau, there have not been any reports from its portion that extends into West Bengal. Hence, our reports probably constitute the eastern most population of this species.

References

- Ball, V., 1874. On the avifauna of the Chutia (Chota) Nagpur Division, S. W. frontier of Bengal. *Stray Feathers* 2 (4&5): 355–440.
- Ball, V., 1878. From the Ganges to the Godavari. On the distribution of birds, so far as it is at present known, throughout the hilly region which extends from the Rajmahal Hills to the Godavari Valley. *Stray Feathers* 7 (3,4&5): 191–235.
- Chakraborty, R., 2011. Mammalian and avian faunal diversity in Damodar Valley under DVC Project Area. *Records of the Zoological Survey of India, Occasional Paper* No. 328: 1–38.
- Grimmett, R., Inskipp, C., & Inskipp, T., 2011. *Birds of the Indian Subcontinent*. 2nd ed. London: Oxford University Press & Christopher Helm. Pp. 1–528.




Fig. 1. Location of the sighting of Painted Spurfowl.

Kumar, P., & Rawat, G. S., 2008. Chotanagpur Plateau: Relict habitats and endemic plants. Pp. 167–173. In: Rawat, G. S., (ed.). 2008. Special habitats and threatened plants of India. *ENVIS Bulletin: Wildlife and Protected Areas* 11 (1): 1–239.

– Supriya Samanta, Debarbab Sen,
Avishek Patra & Saptarshi Mukherjee

Supriya Samanta, Member, Green Plateau, Dept of Zoology, J K College, Purulia West Bengal, India. E-mail: supriyasamanta1234@gmail.com [SS]

Debarbab Sen, Zonal Secretary, Green Plateau, Bishnupur, Bankura, West Bengal, India. E-mail: debarbabsen.bish@gmail.com [DS]

Avishek Patra, Guest Lecturer (Zoology), Bankura Christian College, Bankura, West Bengal, India. E-mail: avishepatra5@gmail.com [AP]

Saptarshi Mukherjee, President, Green Plateau, Mithani, Asansol, West Burdwan, West Bengal, India. E-mail: ahsouhardya@gmail.com [MS]

A leucistic Brown Crake *Zapornia akool* in Karnataka

On the morning of 11 July 2019, while birding with Shyam Hirurkar, we spotted a pair of Brown Crake *Zapornia akool* [164] in Somapur Village (15.444°N, 75.0824°E), Dharwad District, Karnataka. The road we were driving along had a little water channel running parallel to it. We saw a blob of white along the edge of the channel, and initially thought it was domesticated rock pigeon. But as we got closer, it was clear that it was something else. The brown bird followed the partially white one, and we instantly recognised it as a Brown Crake, given its characteristic olive-brown upperparts and greyish face, breast, and belly (Grimmett et al. 2011). The other bird, which accompanied it, was a colour aberrant Brown Crake as parts of its body were white and the rest of it retained the inherent brown colour [165].

A closer examination of the bird in the photographs revealed that it was a partially leucistic individual, as the pattern of white plumage on the bird is bilaterally symmetric and patchy, ruling out any other form of colour aberrations (van Grouw 2013). This is the first record of a leucistic Brown Crake in South Asia (Mahabal et al. 2016).