

Mahanadi River, Odisha, India. *Indian Birds* 13 (1): 1–7.

Shaikh, P. A., Mendis, A., Luis, J., Das, D. K., & Surve, S., 2018. Status and distribution of Indian Skimmer *Rynchops albigollis* in the National Chambal Sanctuary, India. Final Report submitted to BirdLife International. Pp. 1–46.

– Ritvik Singh

House Number 9, Sector-40, Gurgaon, Haryana, India. E-mail: [singhritvik111@gmail.com](mailto:singhritvik111@gmail.com)

### A Brown Boobook *Ninox scutulata* from Nangal, Punjab, India

We report the sighting of a Brown Boobook *Ninox scutulata* from Nangal Wildlife Sanctuary (31.397°N, 76.361°E), Naya Nangal, Rupnagar District, Punjab, India, that occurred on 26 November 2024 at 0845 h. It was perched on a branch of a tree with a thick canopy, in an area with dense undergrowth of *Lantana camara*. Finding an approach for taking good photographs was impossible. However, the best shot that could be managed [163] proved sufficient to identify the species. The tail had broad dark bands and was tipped white. There was a small white patch between the eyes, the upper parts were uniform brown, and the scapulars were white tipped. These identifying features are confirmed in König & Weick (2008) and Taylor (2016).



Paramnoor Singh Antaal

163. Brown Boobook at Nangal Wildlife Sanctuary, Punjab showing broad dark bands on its tail.

In India, the Brown Boobook is distributed widely, throughout the Himalayan foothills, all eastern India, central India, and most of southern India (Rasmussen & Anderton 2012). Until recently, the western extent of its distribution was believed to be the Himalayan foothills of Uttarakhand. However, now the presence of this species has been documented further to the west of Uttarakhand (Abhinav et al. 2023) – from north Haryana, Chandigarh, Himachal Pradesh, and Jammu. Abhinav et al. (2023) also show one old record from Jalalpur, Hoshiarpur District, Punjab, based on a skin in the collection of Frank S. Wright. The date is given as 16 January 1892, and this skin is presently in the collection of Cornell University Museum of Vertebrates (CUMV). We checked the specimen details in GBIF (2025), but the name of the district is not mentioned. We checked with C. Abhinav,

who informed us (in litt., email dated 11 December 2024) that he had chosen the most probable location to depict on the map. We also checked with Vanya Gregor Rohwer, the curator of birds and mammals of CUMV, who informed us (in litt., email dated 08 January 2025) that the specimen is of a female and no district name is indicated in their records. Upon searching for Jalalpur in Punjab on Google Maps, the broader location was suggested as Hoshiarpur district. We also know that there are multiple places by the name Jalalpur in Punjab. We could not find any record of Brown Boobook from other sources, such as journals, social media, and citizen science platforms. Hence, this 1892 record from 'Jalalpur' is the only known record from Punjab until the record presented here. The three closest locations where this species has been recorded are Amb Doli, Pathiar, Kangra district, Himachal Pradesh (Thakur 2023) at a straight-line distance of c.53 km in the northwest direction, Mandi in Himachal Pradesh (Abhinav et al. 2023) at a straight-line distance of c.65 km in the northeast direction, and Chandigarh (Singh 2021; eBird 2025) at a straight-line distance of c.80 km in the southeast direction. Therefore, this new record, coming 132 years after the previous one from Punjab, is significant as it complements other records from neighbouring states that extend the commonly accepted western limit of the species' distribution. It is also likely the first photographic evidence for the state.

### References

- Abhinav, C., Guleria, V., Dogra, P., & Kumar, P., 2023. The range of the Brown Hawk Owl *Ninox scutulata* in northern India—an update. *Indian BIRDS* 19 (1): 13–17.
- eBird, 2025. Species map, Brown Hawk-Owl *Ninox scutulata*. Webpage URL: <https://ebird.org/map/brnhao1>. [Accessed on 26 January 2025].
- GBIF, 2025. CUMV Bird 4343 *Ninox scutulata*. Webpage URL: <https://www.gbif.org/occurrence/1852380621>. [Accessed on 17 August 2025].
- König, C., & Weick, F., 2008. *Owls of the World*. 2<sup>nd</sup> ed. London, United Kingdom: Christopher Helm Publishers. pp. 1–528.
- Singh, V. J., 2021. Rare boobook's noon hoot reveals owl's perch at MCM DAV College. *The Times of India*, Chandigarh, 24 July 2021.
- Taylor, M., 2016. *Owls: A Guide to Every Species in the World*. New York City, United States: Harper Collins. pp. 1–256.
- Thakur, A., 2023. Brown Hawk-Owl, Amb Doli, Pathiar, Kangra district, Himachal Pradesh. Website URL: <https://ebird.org/checklist/S131280607>. [Accessed on 26 January 2025].

– Gurpartap Singh & Paramnoor Singh Antaal

Gurpartap Singh, 1969, Sector 64, Mohali 160062, District Sahibzada Ajit Singh Nagar, Punjab, India. E-mail: [prof.gurpartap.singh@gmail.com](mailto:prof.gurpartap.singh@gmail.com) [GPS] [Corresponding author]  
Paramnoor Singh Antaal, House No. 26, Urban Estate Phase-I, Patiala 147002, Punjab, India. [PSA]

### Two new breeding colonies of the White-rumped Vulture *Gyps bengalensis* in Bangladesh

The Critically Endangered White-rumped Vulture *Gyps bengalensis* was once an abundant and widely distributed raptor in South and South-East Asia (Prakash 1999; Gilbert et al. 2006; Chaudhary et al. 2012; Ghimire et al. 2019), including Bangladesh (Harvey 1990; Thompson & Johnson 1996). These vultures breed colonially or singly on tall trees even occasionally on cliffs, and has a global population estimate of 4,000–6,000 mature individuals (BirdLife International 2025). The species has been declining, especially since the 1990s mainly due to a widely-used painkiller and anti-inflammatory drug diclofenac used for treating livestock (Anonymous 2004; Baral et al. 2005; Cuthbert et al. 2016; BirdLife International 2025). It is the only vulture known to breed regularly in Bangladesh (Khan 2013).

White-rumped Vulture population declined by c.60% in Bangladesh during 2008–2012 (Khan 2013). The last countrywide